

AD-A081 680

NAVY UNDERWATER SOUND LAB NEW LONDON CT
VARIANCE TEST FOR NORMALITY.(U)
DEC 69 J SKORY

F/G 12/1

UNCLASSIFIED

NUSL-TM-2211-343-69

NL

1 of 1
NUSL-TM-2211-343-69




END

DATE

FORMED

4-80

DTIC



ADA081630

✓
LEVEL # 0

(00682)
Copy No. 9
Code No. 21642

(14) NUSL-TM-2211-343-69

NAVY UNDERWATER SOUND LABORATORY
NEW LONDON, CONNECTICUT 06320

(6) VARIANCE TEST FOR NORMALITY.

by

(10) John Skory

USL Problem No.
0-A-055-00-00
ZR 011 01/01

NUSL Technical Memorandum No. 2211-343-69

(11) 30 December 1969

DTIC
ELECTE

INTRODUCTION

For sample sizes up to $n = 50$ an ordered sample $(X_1 \dots X_n)$ may be tested for normality by a calculation involving a vector $(V_1 \dots V_n)$ that is tabulated in (1). If V and X are column vectors the test statistic, also tabulated in (1), is

(12) (3) $W = \frac{(V_1^T X_1)^2}{(X_1 - \bar{X})^2} \quad i = 1 \dots n$

(9) Technical memo

The distribution of W ranges from 0 to 1 where as W approaches 1.0 the distribution of $(X_1 \dots X_n)$ comes closer and closer to being Gaussian.

When dealing with a very large sample, N measurements, how does one proceed to test for normality since V for samples greater than 50 is not available? A common procedure is the chi squared test for deviations from an expected distribution. A more attractive but approximate method, outlined below, is the taking of a sample of $n = 50$ from the cross section of the large sample of N values. These 50 values may

DDG FILE COPY

DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited

254200 Lm

80 1 23 086 Incl 18 to USNUSL Ser 2644-0302

then be tested by $(V_1 \dots V_{50})$ from the table in (1).

METHOD

The procedure for obtaining $n = 50$ values from N values is:

The i^{th} ordered value of the sample $(X_1 \dots X_{50})$
equals the $(NP_i + \frac{50-i}{49})$ th ordered value of sample $(X_1 \dots X_N)$
rounding to the nearest integral value.

The vector $P = (P_1 \dots P_{50})$ is obtained from (2) using 50 percent ranks with the tails of the distribution adjusted outward to give a standard deviation for $(X_1 \dots X_{50})$ close to that of the large sample $(X_1 \dots X_N)$ and with W close to 1. The vector P is:

$P = .0048, .0322, .0531, .0729, .0928, .1126, .1325, .1524, .1722, .1921, .2119,$
 $.2318, .2517, .2715, .2914, .3113, .3311, .3510, .3709, .3907, .4106, .4305,$
 $.4503, .4702, .4901, .5099, .5298, .5497, .5695, .5894, .6093, .6291, .6490,$
 $.6689, .6887, .7086, .7285, .7483, .7682, .7880, .8079, .8278, .8476, .8675,$
 $.8873, .9072, .9270, .9469, .9678, .9952$

If $N = 10,000$ the required set $(X_1 \dots X_{50})$ equals the (49th, 323rd, 532nd ... 9952nd) ordered values of the sample $(X_1 \dots X_{10,000})$.

The vector $(V_1 \dots V_{50}) = (-.3751, \dots -.0035, .0035, \dots, .3751)$ and W would be calculated as usual. From tables of the unit normal distribution a cross-section sample of 50 yielded a sigma of 1.004 (for 50 degrees of freedom) and a W of .9996, both sufficiently close

to unity for general application.

SUMMARY

An alternate method has been outlined to test if a large sample is Gaussian in distribution. Instead of a chi squared test of fit a new statistic W is evaluated using a cross-section sample of 50 from a much larger sample of data. If the large sample is at least 100, the technique yields reliable results which may be assessed for significance against tabulated percentiles of W .

John Skory

JOHN SKORY
Math Statistician

REFERENCES

1. Shapiro, S.S., and Wilk, M.B., "An Analysis of Variance Test for Normality," Biometrika, 52, 3 and 4, p 591, (1965).
2. Failure Distribution Analyses Study, Volume 3, Tables of Plotting Positions, Dept of the Navy, CAI-NY 6052, Contract NObsr-89126, AD631527, (1964).

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>
By <i>Little on file</i>	
Distribution/	
Availability Codes	
Dist.	Availability for special
<i>A</i>	